

## FIRST DETECTION OF MILLIMETER/SUBMILLIMETER EXTRAGALACTIC H<sub>2</sub>O MASER EMISSION

E. M. L. HUMPHREYS, L. J. GREENHILL, M. J. REID, H. BEUTHER, J. M. MORAN, M. GURWELL,  
D. J. WILNER, AND P. T. KONDRATKO

Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138;  
ehumphreys@cfa.harvard.edu, lgreenhill@cfa.harvard.edu, mreid@cfa.harvard.edu,  
hbeuther@cfa.harvard.edu, jmoran@cfa.harvard.edu, mgurwell@cfa.harvard.edu,  
dwilner@cfa.harvard.edu, pkondratko@cfa.harvard.edu

Received 2005 August 18; accepted 2005 October 18; published 2005 November 17

### ABSTRACT

We report the first detection of an extragalactic millimeter wavelength H<sub>2</sub>O maser at 183 GHz toward NGC 3079 using the Submillimeter Array (SMA) and a tentative submillimeter wave detection of the 439 GHz maser toward the same source using the James Clerk Maxwell Telescope (JCMT). These H<sub>2</sub>O transitions are known to exhibit maser emission in star-forming regions and evolved stars. NGC 3079 is a well-studied nuclear H<sub>2</sub>O maser source at 22 GHz with a time-variable peak flux density in the range 3–12 Jy. The 183 GHz H<sub>2</sub>O maser emission, with a peak flux density of ~0.5 Jy (7  $\sigma$  detection), also originates from the nuclear region of NGC 3079 and is spatially coincident with the dust continuum peak at 193 GHz (53 mJy integrated). Peak emission at both 183 and 439 GHz occurs in the same range of velocity as that covered by the 22 GHz spectrum. We estimate the gas-to-dust ratio of the nucleus of NGC 3079 to be  $\approx 150$ , comparable to the Galactic value of 160. Discovery of maser emission in an active galactic nucleus beyond the long-known 22 GHz transition opens the possibility of future position-resolved radiative transfer modeling of accretion disks and outflows <1 pc from massive black holes.

*Subject headings:* galaxies: active — galaxies: individual (NGC 3079) — galaxies: nuclei — masers — submillimeter — techniques: high angular resolution

### 1. INTRODUCTION

Active galactic nuclei (AGNs) that exhibit H<sub>2</sub>O maser emission at 22 GHz have been objects of particular interest since the discovery that the maser emission can trace circumnuclear accretion disks, as in NGC 4258 (Miyoshi et al. 1995; Greenhill et al. 1995a, 1995b). The presence of maser emission enables direct mapping of AGN dynamics within 1 pc of supermassive black holes via very long baseline interferometry (see reviews of Greenhill 2002 and Maloney 2002). The additional discovery that H<sub>2</sub>O maser emission can be associated with shocked gas in jets and outflows has also provided a tracer for a second component in AGNs (Gallimore et al. 1996; Claussen et al. 1998; Peck et al. 2003; Greenhill et al. 2003).

Maser emission at 22 GHz is believed to be predominantly collisionally pumped at gas kinetic temperatures of  $T_k = 400\text{--}1000$  K, hydrogen densities of  $n(\text{H}_2) = 10^8\text{--}10^{10}$  cm<sup>-3</sup>, and dust temperatures  $T_d < 100$  K (e.g., Deguchi 1977; Elitzur et al. 1989; Yates et al. 1997). In AGN accretion disks, gas heating is thought to arise from X-ray irradiation of the disk surfaces, whereas masers in jets and outflows may be heated via shock compressions (Neufeld et al. 1994; Kartje et al. 1999).

Since the physical conditions that give rise to emission at 22 GHz also pump other H<sub>2</sub>O maser transitions, notably those at 183, 321, and 325 GHz (e.g., Deguchi 1977; Neufeld & Melnick 1991; Yates et al. 1997), 22 GHz maser emission is unlikely to be alone in AGN disks, jets, and outflows. We note that, in Galactic regions of star formation and evolved stars, inversion of the 22 GHz transition is accompanied by inversion of other rotational transitions from the ground and vibrationally excited H<sub>2</sub>O states (e.g., Phillips et al. 1980; Waters et al. 1980; Menten & Melnick 1989; Menten et al. 1990a, 1990b; Feldman et al. 1991; Menten & Melnick 1991; Melnick et al. 1993). In AGNs, some of the masers should occur in broadly the same regions as at 22 GHz, but others could probe uncharted regions

of the central engines of AGNs, including regions at smaller radii. Line ratios of two or more maser transitions originating from the same gas would constrain radiative transfer models far better than is now possible. Previous searches for extragalactic millimeter/submillimeter H<sub>2</sub>O masers have been performed but have been hampered by rms sensitivities worse than 1 Jy (K. M. Menten 2004, private communication).

NGC 3079 is an almost edge-on spiral Seyfert 2/LINER galaxy with an optical heliocentric velocity of  $1125 \pm 6$  km s<sup>-1</sup> (e.g., Heckman 1980; Ho et al. 1997), at a distance of 16 Mpc (Sofue 1999). It hosts 22 GHz H<sub>2</sub>O maser emission at velocities between ~940 and 1350 km s<sup>-1</sup>, which are believed to trace a disklike structure at radii of 0.4–1.3 pc from a  $2 \times 10^6 M_\odot$  central engine. The parsec-scale maser disk is aligned with a dense molecular kiloparsec-scale disk (Koda et al. 2002) and is located at the apex of a kiloparsec-scale superbubble (Kondratko et al. 2005; Cecil et al. 2002). The bubble is believed to be inflated by a wide-angle outflow (Cecil et al. 2001) that is also responsible for the radio continuum lobes along the minor axis of the galaxy (e.g., Duric & Seaquist 1988; Trotter et al. 1998; Sawada-Satoh et al. 2000; Yamauchi et al. 2004). Within a parsec of the nucleus, the relativistic jet appears aligned with the inner wall of the bubble and may be responsible for weak 22 GHz maser emission away from the disk (Kondratko et al. 2005 and references therein). All of these emission sources lie within the primary beam of the SMA observations reported here.

### 2. OBSERVATIONS AND DATA REDUCTION

#### 2.1. SMA Observations at 183 GHz

We observed the ground-state  $3_{13} \rightarrow 2_{20}$  transition of para-H<sub>2</sub>O ( $E_u/k \sim 200$  K) at a rest frequency of 183.310 GHz ( $\lambda =$

1.6 mm) toward NGC 3079 using the SMA<sup>1</sup> on 2005 March 1. The compact configuration included six antennas, resulting in projected baselines of 8–72 m. The SMA receivers operate in double-sideband mode with two 2 GHz sidebands that we tuned to 183.310 and 193.310 GHz (band center). The correlator provided continuous coverage of each sideband (in 24 separate chunks of 256 channels) and 0.4 MHz channel spacing. The phase center of the observations was  $\alpha_{2000} = 10^{\text{h}}01^{\text{m}}58^{\text{s}}.53$ ,  $\delta_{2000} = 55^{\circ}40'50''.1$  (Cotton et al. 1999). The zenith opacity measured with the NRAO tipping radiometer located at the Caltech Submillimeter Observatory was  $\tau(225 \text{ GHz}) = 0.03\text{--}0.05$  throughout.

We calibrated the data within the IDL superset MIR developed for the Owens Valley Radio Observatory (Scoville et al. 1993) and adapted for the SMA; the imaging was performed in MIRIAD. Data reduction took into account the effect of the pressure-broadened terrestrial H<sub>2</sub>O absorption line centered at 183.310 GHz. Fortunately, the redshift of the galaxy reduces the line frequency of the maser transition by  $\sim 0.7$  GHz, greatly lowering the corresponding atmospheric opacity. For bandpass calibration, we used Jupiter and Saturn. Phase calibration was performed via frequent observations of the quasar 0923+392 about  $18^{\circ}$  from the phase center. We performed the amplitude calibration separately for upper and lower sideband data. For the upper sideband, we used 0923+392; for lower sideband data, we performed amplitude calibration on each of the 24 spectral chunks separately using Saturn and Jupiter. The calibration of the lower sideband as a function of both elevation and frequency was designed to reduce the effect of variation in terrestrial H<sub>2</sub>O absorption across the sideband. We derived the flux scale using observations of Callisto with an estimated accuracy of 20%. Single-sideband system temperatures corrected to the top of the atmosphere were between 400 and 850 K.

## 2.2. JCMT Observations at 439 GHz

We observed the ground-state  $6_{43} \rightarrow 5_{50}$  transition of ortho-H<sub>2</sub>O ( $E_u/k \sim 1100$  K) at a rest frequency of 439.151 GHz ( $\lambda = 0.7$  mm) toward NGC 3079 using the 15 m JCMT on 2004 November 18 in  $\tau_{22.5} = 0.041\text{--}0.045$  weather. The instantaneous bandwidth was 1840 MHz, corresponding to  $\sim 1200$  km s<sup>-1</sup>, with a channel spacing of 1250 kHz or 0.85 km s<sup>-1</sup>. The integration time on-source was 2 hr in beam-switching mode with a throw of  $60''$ . The average  $T_{\text{sys}}$  during the observations was 2650 K. We reduced the data using the Starlink spectral line reduction software, SpecX.

## 3. RESULTS

We detected millimeter (193 GHz) continuum emission with a peak signal-to-noise ratio (S/N) of  $\sim 10$  (Fig. 1) at  $\alpha_{2000} = 10^{\text{h}}01^{\text{m}}57^{\text{s}}.80 \pm 0^{\text{s}}.02$ ,  $\delta_{2000} = 55^{\circ}40'46''.9 \pm 0''.3$  (statistical uncertainties). Systematic uncertainties of (0''.02, 0''.3) were estimated through measurement of an apparent position for 1150+497 against the phase calibrator 0923+392. These place the peak of the millimeter continuum within  $1 \sigma$  of the 22 GHz maser position. We partially resolved the nucleus in the north-south direction, and the elongation is approximately parallel to the position angles of both the parsec-scale accretion disk traced by 22 GHz maser emission and the kiloparsec-scale molecular disk detected using, e.g., CO (Koda et al. 2002).

<sup>1</sup> The SMA is a joint project between the Smithsonian Astrophysical Observatory and the Academia Sinica Institute of Astronomy and Astrophysics, and is funded by the Smithsonian Institution and the Academia Sinica.




FIG. 1.—*Filled color contours*: Millimeter continuum image of NGC 3079 at 193 GHz observed with the SMA. The peak flux density is  $27 \text{ mJy beam}^{-1}$  with a  $1 \sigma$  rms of  $2.5 \text{ mJy beam}^{-1}$ ; the integrated continuum flux density is  $53 \text{ mJy}$ . *Line contours*: 183 GHz maser emission at  $1017 \text{ km s}^{-1}$  (optical, heliocentric). The peak maser flux density in the image is  $0.39 \text{ Jy beam}^{-1}$ . Positive emission is shown as solid lines, and negative features as dashed contours. Contours are  $k \times 2 \sigma$ , where  $k = -1, 1, 2, 3$ , and  $1 \sigma = 56 \text{ mJy beam}^{-1}$ . The black cross marks the position of peak 22 GHz H<sub>2</sub>O maser emission at  $956 \text{ km s}^{-1}$ :  $\alpha_{2000} = 10^{\text{h}}01^{\text{m}}57^{\text{s}}.802 \pm 0^{\text{s}}.001$ ,  $\delta_{2000} = 55^{\circ}40'47''.26 \pm 0''.01$  (Kondratko et al. 2005). The black solid arrow indicates the position angles (P.A.'s;  $\sim 10^{\circ}$  east of north) of the parsec-scale 22 GHz H<sub>2</sub>O maser disk (Kondratko et al. 2005) and of the kiloparsec-scale CO disk (Koda et al. 2002). The black dotted arrow marks the direction of relativistic outflow, with the uncertainty depicted by bracketing solid lines of P.A. =  $126^{\circ} \pm 5^{\circ}$  (Trotter et al. 1998; Sawada-Satoh et al. 2000; Middelberg et al. 2004; Kondratko et al. 2005). The arc to the east denotes the opening angle of the kiloparsec-scale superbubble outflowing from the galactic nucleus (e.g., Cecil et al. 2002). The SMA synthesized beam has an FWHM of  $5''.9 \times 3''.4$  (upper sideband) and  $6''.1 \times 3''.6$  (lower sideband, shown top left), with a P.A. of  $27^{\circ}$  in both cases.

We detected 183 GHz H<sub>2</sub>O emission with a peak S/N of  $\sim 7$  in both the real part of the amplitude spectrum and the interferometer phase spectrum (Figs. 1 and 2). The peak flux density is  $0.55 \text{ Jy}$  at  $1017 \text{ km s}^{-1}$ . The line covers nine frequency channels and overlaps a persistent Doppler component appearing in published 22 GHz maser spectra for dates 1998 May 8, 2000 March 18 (where it is especially prominent at  $\sim 2.5 \text{ Jy}$ ; Hagiwara et al. 2002), 2001 March 23 (Kondratko et al. 2005), and 2002 April 12 (Braatz et al. 2003). We note the possible detection of additional spectral features at 1208 (five channels wide) and  $1265 \text{ km s}^{-1}$  (four channels wide). Both features appear in spectra of real amplitude and phase (near-zero degrees, which indicates positions similar to the  $1017 \text{ km s}^{-1}$  emission), and both lie within the velocity interval of 22 GHz emission. Note that the correspondence of features between 22 and 183 GHz observations is not precise and is also not expected to be true for all features under certain pumping scenarios. The peak maser emission is at  $\alpha_{2000} = 10^{\text{h}}01^{\text{m}}57^{\text{s}}.75 \pm 0^{\text{s}}.03$ ,  $\delta_{2000} = 55^{\circ}40'46''.7 \pm 0''.4$ , which is within  $1 \sigma$  of the 193 GHz continuum peak. At our resolution, we cannot determine whether 183 GHz emission originates from the disk or outflow, or from both. We note that in addition to tracing a disk, a subset of 22 GHz maser emission is associated with the outflow within a few parsecs of the nucleus (Kondratko et al. 2005). The 22 GHz emission at around 1000 and  $1200 \text{ km s}^{-1}$  is present in both the disk and outflow. Possible angular extension of the 183 GHz emission to the southeast is marginally detected ( $2 \sigma$ ), which lies along the axis of the relativistic flow (e.g., Kondratko et al. 2005). We note that Hagiwara et al.




FIG. 2.—The 183 GHz maser spectrum toward the nucleus of NGC 3079 corrected for atmospheric absorption obtained with the SMA. *Top panel*: Real part of the spectrum exhibiting an  $\sim 0.55$  Jy feature at a velocity of  $1017 \text{ km s}^{-1}$ . The channel spacing is  $5 \text{ km s}^{-1}$ . The atmospheric opacity for a zenith angle of  $50^\circ$  is shown by the dashed line; peaks at  $1850$  and  $2150 \text{ km s}^{-1}$  are due to ozone. Note that the noise rises at lower velocities due to the effect of atmospheric absorption. *Upper bottom panel*: 22 GHz spectrum obtained with a Deep Space Network (DSN) 70 m antenna (P. T. Kondratko 2005, private communication) between  $800$  and  $1900 \text{ km s}^{-1}$ . *Middle bottom panel*: Real part of the 183 GHz SMA spectrum between  $800$  and  $1900 \text{ km s}^{-1}$ . *Lower bottom panel*: Corresponding fringe phase. Tentative maser features at  $1208$  and  $1265 \text{ km s}^{-1}$  are also noted with arrows.

(2004) detect OH absorption associated with the outflow at an overlapping velocity and that we cannot rule out shocks in circumnuclear material as an origin for 183 GHz emission.

We obtained a tentative detection ( $5 \sigma$ ) of the 439 GHz maser at  $1157 \text{ km s}^{-1}$  (Fig. 3). With peak  $S_\nu \sim 2.5 \text{ Jy}$ , 439 GHz emission is of comparable strength to typical values for peak 22 GHz emission toward NGC 3079 and occurs within the 22 GHz velocity interval.

## 4. DISCUSSION

### 4.1. Millimeter Dust Emission

The millimeter continuum of NGC 3079 is probably dominated by thermal emission from optically thin dust. As in the  $1.2 \text{ mm}$  observations of Braine et al. (1997) at  $11''$  resolution, we find dust concentrated in the nuclear region of NGC 3079 (Fig. 1) and place an upper limit of  $8 \text{ mJy}$  ( $3 \sigma$ ) on emission associated with the radio lobes  $\sim 1 \text{ kpc}$  from the emission peak. Braine et al. (1997) estimate the dust temperature to be  $\sim 30 \text{ K}$  at kiloparsec radii. However, in two other AGNs, the presence of hotter dust at 100 times smaller radii, of  $\sim 300 \text{ K}$ , has been inferred from VLT observations (Jaffe et al. 2004; Prieto et al. 2004). Adopting a dust temperature of  $100 \text{ K}$  for NGC 3079 on intermediate scales and a dust opacity index  $\beta = 2$ , we estimate that the enclosed dust mass of the inner  $200 \text{ pc}$  is  $M_{\text{dust}} \sim 10^{6.3 \pm 0.8} M_\odot$  using the method described in Hildebrand (1983). Using CO Nobeyama observations by Koda et al. (2002), who find a gas mass of  $M_{\text{gas}} \sim 10^{8.5} M_\odot$  within the central  $150 \text{ pc}$ , we estimate a gas-to-dust ratio of 150, which is comparable to the Galactic value of 160 and to the ratios derived in bright local galaxies (Dunne et al. 2000).

### 4.2. The 183 GHz Emission

The 183 GHz emission that we detect in the nucleus of NGC 3079 is likely from a maser process. It arises from one, and possibly more, narrow unresolved features in the same range of velocity as that of known 22 GHz maser emission. From radiative transfer models, we know that the same high densities and temperature conditions that produce maser emission at 22 GHz also strongly invert the 183 GHz H<sub>2</sub>O transition. Noting




FIG. 3.—*Top*: 22 GHz spectrum obtained using the Green Bank Telescope (GBT) from Braatz et al. (2003). *Bottom*: 439 GHz H<sub>2</sub>O emission toward NGC 3079 observed with the JCMT. The peak flux density is  $2.5 \text{ Jy}$ , and the  $1 \sigma$  rms is  $0.5 \text{ Jy}$ . The channel spacing is  $13.7 \text{ km s}^{-1}$ .

that the 22 GHz feature at  $1017 \text{ km s}^{-1}$  varies between 0.4 and 2.6 Jy (Hagiwara et al. 2002), and assuming an ortho : para  $\text{H}_2\text{O}$  ratio of 3 : 1, pumping models could reproduce likely line ratios (e.g., Yates et al. 1997). Narrow line widths ( $<8 \text{ km s}^{-1}$ ) at 183 GHz, small compared with the characteristic rotational and apparent random velocities for molecular gas in this nucleus (Kondratko et al. 2005), also indicate that emitting regions should be compact. Our beam size is too large to determine whether emission is thermal or maser based on brightness temperature arguments. For the emission to be thermal, with a brightness temperature of  $\leq 2000 \text{ K}$  such that  $\text{H}_2\text{O}$  molecules are not dissociated, the emitting region must be  $>0''.1$  (10 pc).

#### 4.3. The 439 GHz Emission

The 439 GHz emission is probably also due to maser amplification. Emission occurs over a narrow velocity range (Fig. 3), and conditions for strong 439 GHz inversion are likely to exist in the galactic nucleus. The 22 GHz spectrum peaks to the blue of the  $v_{\text{sys}}$  at  $1125 \text{ km s}^{-1}$ , yet the 439 GHz spectrum peaks slightly above  $v_{\text{sys}}$  at  $1157 \text{ km s}^{-1}$  (Fig. 3). We ascribe this difference to the different pump mechanisms operating for each maser. Whereas 22 GHz emission is collisionally pumped and

is increasingly quenched by local dust temperatures  $>100 \text{ K}$ , the 439 GHz maser is predominantly radiatively pumped and is most strongly inverted in the presence of  $T_d = 300 \text{ K}$ ,  $T_k = 400 \text{ K}$ , and  $n(\text{H}_2) = 5 \times 10^9 \text{ cm}^{-3}$  (Yates et al. 1997). Dust temperatures of this order and higher are reasonable in the central parsecs of the nucleus (Jaffe et al. 2004; Prieto et al. 2004). Peak 439 GHz emission ( $1157 \text{ km s}^{-1}$ ) is therefore likely to originate from a hot, dusty region around the central engine, possibly at radii  $\ll 1 \text{ pc}$ , separated spatially from the gas emitting most strongly at 22 GHz ( $956 \text{ km s}^{-1}$ ). At 22 GHz, features at these velocities are indeed separated spatially by  $\sim 1 \text{ pc}$  projected (Kondratko et al. 2005). The observations in Figure 3 were taken on different dates, and we note that the 22 GHz feature near  $1157 \text{ km s}^{-1}$  is not present at all epochs (e.g., in Fig. 2). However, the feature line ratio in Figure 3 of  $\sim 100$  is accommodated by the vicinity of hot dust (Yates et al. 1997).

We thank all the staff at SMA and JCMT. We thank S. Paine for opacity data, J. Braatz for 22 GHz data, and G. Melnick for useful discussions. H. B. acknowledges support from the Emmy-Noether Program of the Deutsche Forschungsgemeinschaft (DFG-BE2578/1).

#### REFERENCES

- Braatz, J. A., Wilson, A. S., Henkel, C., Gough, R., & Sinclair, M. 2003, *ApJS*, 146, 249
- Braine, J., Guélin, M., Dumke, M., Brouillet, N., Herpin, F., & Wielebinski, R. 1997, *A&A*, 326, 963
- Cecil, G., Bland-Hawthorn, J., & Veilleux, S. 2002, *ApJ*, 576, 745
- Cecil, G., Bland-Hawthorn, J., Veilleux, S., & Filippenko, A. V. 2001, *ApJ*, 555, 338
- Claussen, M. J., Diamond, P. J., Braatz, J. A., Wilson, A. S., & Henkel, C. 1998, *ApJ*, 500, L129
- Cotton, W. D., Condon, J. J., & Arbizzani, E. 1999, *ApJS*, 125, 409
- Deguchi, S. 1977, *PASJ*, 29, 669
- Dunne, L., Eales, S., Edmunds, M., Ivison, R., Alexander, P., & Clements, D. L. 2000, *MNRAS*, 315, 115
- Duric, N., & Seaquist, E. R. 1988, *ApJ*, 326, 574
- Elitzur, M., Hollenbach, D. J., & McKee, C. F. 1989, *ApJ*, 346, 983
- Feldman, P. A., Matthews, H. E., Cunningham, C. T., Hayward, R. H., Wade, J. D., Amano, T., & Scappini, F. 1991, *JRASC*, 85, 191
- Gallimore, J. F., Baum, S. A., O'Dea, C. P., Brinks, E., & Pedlar, A. 1996, *ApJ*, 462, 740
- Greenhill, L. 2002, in *IAU Symp. 206, Cosmic Masers: From Proto-Stars to Black Holes*, ed. V. Migenes & M. J. Reid (San Francisco: ASP), 381
- Greenhill, L. J., Henkel, C., Becker, R., Wilson, T. L., & Wouterloot, J. G. A. 1995a, *A&A*, 304, 21
- Greenhill, L. J., Jiang, D. R., Moran, J. M., Reid, M. J., Lo, K. Y., & Claussen, M. J. 1995b, *ApJ*, 440, 619
- Greenhill, L. J., et al. 2003, *ApJ*, 590, 162
- Hagiwara, Y., Henkel, C., Sherwood, W. A., & Baan, W. A. 2002, *A&A*, 387, L29
- Hagiwara, Y., Klöckner, H.-R., & Baan, W. A. 2004, *MNRAS*, 353, 1055
- Heckman, T. M. 1980, *A&A*, 87, 152
- Hildebrand, R. H. 1983, *QJRAS*, 24, 267
- Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1997, *ApJS*, 112, 315
- Jaffe, W., et al. 2004, *Nature*, 429, 47
- Kartje, J. F., Königl, A., & Elitzur, M. 1999, *ApJ*, 513, 180
- Koda, J., Sofue, Y., Kohno, K., Nakanishi, H., Onodera, S., Okumura, S. K., & Irwin, J. A. 2002, *ApJ*, 573, 105
- Kondratko, P. T., Greenhill, L. J., & Moran, J. M. 2005, *ApJ*, 618, 618
- Maloney, P. R. 2002, *Publ. Astron. Soc. Australia*, 19, 401
- Melnick, G. J., Menten, K. M., Phillips, T. G., & Hunter, T. 1993, *ApJ*, 416, L37
- Menten, K. M., & Melnick, G. J. 1989, *ApJ*, 341, L91
- . 1991, *ApJ*, 377, 647
- Menten, K. M., Melnick, G. J., & Phillips, T. G. 1990a, *ApJ*, 350, L41
- Menten, K. M., Melnick, G. J., Phillips, T. G., & Neufeld, D. A. 1990b, *ApJ*, 363, L27
- Middelberg, E., et al. 2004, *A&A*, 417, 925
- Miyoshi, M., Moran, J., Herrnstein, J., Greenhill, L., Nakai, N., Diamond, P., & Inoue, M. 1995, *Nature*, 373, 127
- Neufeld, D. A., Maloney, P. R., & Conger, S. 1994, *ApJ*, 436, L127
- Neufeld, D. A., & Melnick, G. J. 1991, *ApJ*, 368, 215
- Peck, A. B., Henkel, C., Ulvestad, J. S., Brunthaler, A., Falcke, H., Elitzur, M., Menten, K. M., & Gallimore, J. F. 2003, *ApJ*, 590, 149
- Phillips, T. G., Kwan, J., & Huggins, P. J. 1980, in *IAU Symp. 87, Interstellar Molecules*, ed. B. H. Andrew (Dordrecht: Reidel), 21
- Prieto, M. A., et al. 2004, *ApJ*, 614, 135
- Sawada-Satoh, S., Inoue, M., Shibata, K. M., Kamenoi, S., Migenes, V., Nakai, N., & Diamond, P. J. 2000, *PASJ*, 52, 421
- Scoville, N. Z., Carlstrom, J. E., Chandler, C. J., Phillips, J. A., Scott, S. L., Tilanus, R. P. J., & Wang, Z. 1993, *PASP*, 105, 1482
- Sofue, Y. 1999, *Adv. Space Res.*, 23, 949
- Trotter, A. S., Greenhill, L. J., Moran, J. M., Reid, M. J., Irwin, J. A., & Lo, K. 1998, *ApJ*, 495, 740
- Waters, J. W., et al. 1980, *ApJ*, 235, 57
- Yamauchi, A., Nakai, N., Sato, N., & Diamond, P. 2004, *PASJ*, 56, 605
- Yates, J. A., Field, D., & Gray, M. D. 1997, *MNRAS*, 285, 303